

Women Proudly Caring for Women

275 Collier Road NW Suite 230 Atlanta, GA 30309 www.awog.org

Telephone(404Facsimile(404Answering Service(404

(404) 352-3616 (404) 352-2028 (404) 487-2134

Nausea and Vomiting Diet

- 1. Eat every 2-3 hrs, small amounts.
- 2. Do not eat and drink at the same time.
- 3. Take your prenatal vitamin at night, right before bed.
- 4. Try Ginger Tea, Canned Ginger, or real ginger ale.
- 5. Try Starchy foods: Rice, Pasta, Potatoes, Breads, and Crackers.
- 6. Hydration is most important- Water, Gatorade, Ice Chips, Jello, Popsicles
- 7. Try Peppermint candies or tea.
- 8. Try Lemons and lemon drops, can help if you have continuous saliva.
- 9. Try to eat some protein throughout day-Peanut butter, broiled chicken, soy or dairy sources, nuts, canned fruits, eggs if tolerated.
- 10.Try canned peaches in HEAVY syrup-if not diabetic-sip chilled liquid 1-2 tablespoons at a time- try hydrating after 15 min.
- 11.Get fresh air during the day.
- 12. "Sea Bands" motion sickness wrist bands from any pharmacy.
- 13. Take 25mg Pyridoxine (Vitamin B6) & 12.5mg Doxylamine (Unisom)

*Usually resolves around 12 weeks.

- * Sometimes medication may be needed
- *Call our office if you feel it is more than morning sickness